

# National Air Quality and Emissions Trends Report, 1999

U.S. Environmental Protection Agency  
Office of Air Quality Planning and Standards  
Emissions Monitoring and Analysis Division  
Air Quality Trends Analysis Group  
Research Triangle Park, North Carolina 27711

March 2001

### **About the Cover**

The map on the cover depicts nationwide annual mean PM<sub>2.5</sub> concentrations from the Federal Reference Method (FRM) monitoring network, as well as information on data completeness. Annual mean concentrations are generally above the level of the 1997 standard of 15 µg/m<sup>3</sup> in much of the eastern United States and throughout California. Annual mean concentrations above 20 µg/m<sup>3</sup> are seen in several major metropolitan areas including Pittsburgh, Cleveland, Atlanta, Chicago, and St. Louis and Los Angeles. The western Great Plains and mountain regions show notably low annual mean concentrations, most below 10 µg/m<sup>3</sup>.

Data Source: U.S. EPA AIRS Data Base 1/30/01.

### **Disclaimer**

This report has been reviewed and approved for publication by the U.S. Environmental Protection Agency's Office of Air Quality Planning and Standards. Mention of trade names or commercial products are not intended to constitute endorsement or recommendation for use.

### **Acknowledgments**

The Trends Team would like to acknowledge the members of EPA's Office of Research and Development, Office of Atmospheric Programs, Office of Radiation and Indoor Air and Office of Transportation and Air Quality for peer reviewing this report prior to publication; Support for the statistical analyses of air toxics trends provided under EPA contract 68D70066; Colorado State University for providing summary data from the IMPROVE monitoring network; Support for desktop publishing and Web site development provided under EPA contract 68W99004; and the Trends Workgroup in EPA's Office of Air Quality Planning and Standards for providing comments throughout report development.

## **Preface**

This is the 27th annual report on air pollution trends in the United States issued by the U.S. Environmental Protection Agency. The report is prepared by the Air Quality Trends Analysis Group (AQTAG) in Research Triangle Park, North Carolina and is directed toward both the technical air pollution audience and other interested parties and individuals.

The report can be accessed via the Internet at <http://www.epa.gov/airtrends/>. AQTAG solicits comments on this report and welcomes suggestions regarding techniques, interpretations, conclusions, or methods of presentation. Comments can be submitted via the website or mailed to:

Attn: Trends Team  
AQTAG (MD-14)  
U.S. EPA  
Research Triangle Park, NC 27711

Readers can access data from the Aerometric Information Retrieval System (AIRS) at <http://www.epa.gov/airsdata/> and real time air pollution data at <http://www.epa.gov/airnow/>.



# Contents

## Chapter 1

<b>Executive Summary .....</b>	<b>1</b>
Chapter 2 .....	2
Chapter 3 .....	2
Chapter 4 .....	3
Chapter 5 .....	3
Chapter 6 .....	4
Chapter 7 .....	6
References and Notes .....	7

## Chapter 2

<b>Criteria Pollutants — National Trends .....</b>	<b>9</b>
Carbon Monoxide .....	11
Lead .....	17
Nitrogen Dioxide .....	23
Ozone .....	29
Particulate Matter .....	40
Sulfur Dioxide .....	61
References .....	67

## Chapter 3

<b>Criteria Pollutants — Metropolitan Area Trends.....</b>	<b>69</b>
Status: 1999 .....	69
Trends Analysis .....	70
The Air Quality Index .....	70
Summary of AQI Analyses .....	72
References and Notes .....	74

## Chapter 4

<b>Criteria Pollutants — Nonattainment Areas .....</b>	<b>75</b>
--	-----------

<b>Chapter 5</b>	
Air Toxics .....	<b>79</b>
Background .....	79
Health and Environmental Effects .....	79
National Air Toxics Control Program .....	80
Air Toxics Emissions in 1996 .....	82
Trends in Air Toxics Emissions .....	84
Ambient Monitoring .....	84
Trends In Ambient Concentrations .....	85
National Atmospheric Deposition Program/Mercury Deposition Network .....	96
Integrated Atmospheric Deposition Network .....	96
References .....	97
<b>Chapter 6</b>	
Visibility Trends .....	<b>99</b>
Introduction .....	99
Nature and Sources of the Problem .....	100
Long-Term Trends (1981–1995) .....	102
Recent Trends (1990–1999) from IMPROVE Data .....	103
Regional Visibility Trends for the Eastern and Western United States .....	104
The Components of PM Contributing to Trends in Visibility Impairment .....	105
Current Visibility Conditions .....	109
Programs to Improve Visibility .....	113
References .....	114
<b>Chapter 7</b>	
Atmospheric Deposition of Sulfur and Nitrogen Compounds .....	<b>115</b>
Primary Atmospheric Deposition Monitoring Networks .....	116
National Atmospheric Deposition Network/National Trends Network .....	116
Trends Analyses for Sulfate and Nitrogen Concentrations in Wet Deposition .....	117
Clean Air Status and Trends Network .....	118
Dry Deposition .....	119
Concentration Trends Analysis at CASTNet Sites .....	119
Seasonal Trends in SO <sub>2</sub> Emissions and Related Air Quality .....	122
Sulfur and Nitrogen Deposition .....	124
References .....	125
<b>Appendix A</b>	
Data Tables .....	<b>127</b>
<b>Appendix B</b>	
Methodology .....	<b>229</b>
AIRS Methodology .....	229
Emissions Estimates Methodology .....	233
IMPROVE Methodology .....	234
Air Toxics Methodology .....	234
References .....	237

# Figures

Figure 2-1.	Trend in 2nd maximum non-overlapping 8-hour average CO concentrations, 1980–1999. ....	11
Figure 2-2.	Trend in 2nd maximum non-overlapping 8-hour average CO concentrations by type of location, 1980–1999. ....	12
Figure 2-3.	Trend in CO 2nd maximum non-overlapping 8-hour concentrations by EPA region, 1980–1999. ....	12
Figure 2-4.	CO emissions by source category, 1999. ....	13
Figure 2-5.	Density map of 1999 carbon monoxide emissions, by county. ....	13
Figure 2-6.	Trend in national total CO emissions, 1980–1999. ....	14
Figure 2-7.	Highest 2nd maximum non-overlapping 8-hour average CO concentration by county, 1999. ....	15
Figure 2-8.	Trend in maximum quarterly average Pb concentrations (excluding point-source oriented sites), 1980–1999. ....	18
Figure 2-9.	Pb maximum quarterly mean concentration trends by location (excluding point-source oriented sites), 1980–1999. ....	18
Figure 2-10.	Trend in Pb maximum quarterly mean concentration by EPA region, 1980–1999. ....	19
Figure 2-11.	National total Pb emissions trend, 1980–1999. ....	20
Figure 2-12.	Pb emissions by source category, 1999. ....	20
Figure 2-13.	Pb maximum quarterly concentration in the vicinity of Pb point sources, 1999. ....	21
Figure 2-14.	Highest Pb maximum quarterly mean by county, 1999. ....	21
Figure 2-15.	Trend in annual NO <sub>2</sub> mean concentrations, 1980–1999. ....	24
Figure 2-16.	Trend in annual mean NO <sub>2</sub> concentrations by type of location, 1980–1999. ....	24
Figure 2-17.	Trend in NO <sub>2</sub> maximum quarterly mean concentration by EPA region, 1980–1999. ....	25
Figure 2-18.	Trend in national total NO <sub>x</sub> emissions, 1980–1999. ....	26
Figure 2-19.	NO <sub>x</sub> emissions by source category, 1999. ....	26
Figure 2-20.	Density map of 1999 nitrogen dioxide emissions, by county. ....	27
Figure 2-21.	Highest NO <sub>2</sub> annual mean concentration by county, 1999. ....	28
Figure 2-22.	Trend in annual 2nd-highest daily maximum 1-hour, and 4th-highest daily 8-hour O <sub>3</sub> concentrations, 1980–1999. ....	30
Figure 2-23.	Trend in 4th-highest daily 8-hour O <sub>3</sub> concentrations, 1980–1999. ....	31
Figure 2-24.	Trend in 2nd highest daily 1-hour O <sub>3</sub> concentration by EPA region, 1980–1999. ....	32
Figure 2-25.	Trend in 4th highest daily 8-hour O <sub>3</sub> concentration by EPA region, 1980–1999. ....	32
Figure 2-26.	Trend in annual 2nd-highest daily maximum 1-hour O <sub>3</sub> concentrations by location, 1980–1999. ....	33
Figure 2-27.	Comparison of actual and meteorologically adjusted 1-hour O <sub>3</sub> trends, 1980–1999. ....	33
Figure 2-28.	Areas with PAMS networks. ....	34
Figure 2-29.	A comparison of the median change in summer morning concentrations of the most abundant VOC species measured at all PAMS sites and PAMS type 2 sites from 1995 and 1999. ....	34
Figure 2-30.	Trend in 4th-highest daily 8-hour O <sub>3</sub> based on 34 CASTNet sites in the rural eastern United States, 1980–1999. ....	35
Figure 2-31.	Trend in annual 4th-highest daily maximum 8-hour O <sub>3</sub> concentrations in National Parks, 1980–1999. ....	35
Figure 2-32.	Trend in national total anthropogenic VOC emissions, 1980–1999. ....	36
Figure 2-33.	Anthropogenic VOC emissions by source category, 1999. ....	36
Figure 2-34.	Density map of 1999 VOC emissions, by county. ....	38
Figure 2-35.	Highest second daily maximum 1-hour O <sub>3</sub> concentration by county, 1999. ....	38
Figure 2-36.	Highest fourth daily maximum 8-hour O <sub>3</sub> concentration by county, 1999. ....	39
Figure 2-37.	Trend in annual mean PM <sub>10</sub> concentrations, 1990–1999. ....	42
Figure 2-38.	PM <sub>10</sub> annual mean concentration trends by location, 1990–1999. ....	42
Figure 2-39.	National PM <sub>10</sub> emissions trend, 1980–1999 (traditionally inventoried sources only). ....	43
Figure 2-40.	PM <sub>10</sub> emissions from traditionally inventoried source categories, 1999. ....	43

Figure 2-41.	Total PM <sub>10</sub> emissions by source category, 1999 .....	44
Figure 2-42.	PM <sub>10</sub> emissions density by county, 1999.....	45
Figure 2-43.	Trend in PM <sub>10</sub> annual mean concentration by EPA region, 1990–1999. ....	45
Figure 2-44.	Highest 2nd maximum 24-hour PM <sub>10</sub> concentration by county, 1999. ....	46
Figure 2-45.	Status of PM <sub>2.5</sub> monitor network, as of May 2001. ....	47
Figure 2-46.	1999 annual mean PM <sub>2.5</sub> concentrations (µg/m <sup>3</sup> ). ....	48
Figure 2-47.	1999 98th percentile 24-hour average PM <sub>2.5</sub> concentrations (µg/m <sup>3</sup> ). ....	48
Figure 2-48.	Urban PM <sub>2.5</sub> monthly patterns by region, 1999. ....	49
Figure 2-49.	Rural PM <sub>2.5</sub> monthly patterns by region, 1999. ....	50
Figure 2-50.	Class I Areas in the IMPROVE Network meeting the data completeness criteria in Appendix B. ....	51
Figure 2-51.	Annual mean PM <sub>2.5</sub> concentrations in 1999. ....	52
Figure 2-52.	PM <sub>2.5</sub> concentrations, 1992–1999 at eastern IMPROVE sites meeting trends criteria. ....	52
Figure 2-53.	PM <sub>2.5</sub> concentrations, 1990–1999 at western IMPROVE sites meeting trends criteria. ....	53
Figure 2-54.	PM <sub>2.5</sub> concentrations, 1990–1999 at the Washington D.C. IMPROVE site. ....	53
Figure 2-55.	PM <sub>2.5</sub> ambient composition. ....	55
Figure 2-56.	PM <sub>2.5</sub> emission sources. ....	56
Figure 2-57.	Direct PM <sub>2.5</sub> emissions density by county, 1999. ....	57
Figure 2-58.	National direct PM <sub>2.5</sub> emissions trend, 1990–1999 (traditionally inventoried sources only). ....	58
Figure 2-59.	Direct PM <sub>2.5</sub> emissions from traditionally inventoried source categories, 1999. ....	58
Figure 2-60.	Total direct PM <sub>2.5</sub> emissions by source category, 1999. ....	59
Figure 2-61.	National ammonia emissions by principle source categories, 1999. ....	59
Figure 2-62.	Estimated 1999 annual mean PM <sub>10</sub> -2.5. ....	60
Figure 2-63.	Estimated 1999 98th percentile 24-hour average PM <sub>10</sub> -2.5 developed from 1999 FRM monitor data. ....	60
Figure 2-64.	Trend in annual mean SO <sub>2</sub> concentrations, 1980–1999. ....	62
Figure 2-65.	Annual mean SO <sub>2</sub> concentration by trend location, 1980–1999. ....	62
Figure 2-66.	National total SO <sub>2</sub> emissions trend, 1980–1999. ....	63
Figure 2-67.	SO <sub>2</sub> emissions by source category, 1999. ....	63
Figure 2-68.	Long-term ambient SO <sub>2</sub> trend, 1980–1999. ....	64
Figure 2-69.	Trend in SO <sub>2</sub> annual arithmetic mean concentration by EPA region, 1980–1999. ....	65
Figure 2-70.	Plants affected by the Acid Rain Program. ....	65
Figure 2-71.	Highest 2nd maximum 24-hour SO <sub>2</sub> concentration by county, 1999. ....	66
Figure 3-1.	Air Quality Index logo. ....	72
Figure 3-2.	Number of days with AQI values > 100, as a percentage of 1990 value. ....	73
Figure 3-3.	Percent of days over 100 due to ozone. ....	73
Figure 4-1.	Location of nonattainment areas for criteria pollutants, September 2000. ....	75
Figure 4-2.	Classified ozone nonattainment areas. ....	76
Figure 5-1.	National contribution of source types to 1996 NTI emissions for the 188 HAPs. ....	82
Figure 5-2.	National contribution of source types to 1996 NTI emissions for the urban HAPs. ....	82
Figure 5-3.	National contribution by emission source type for individual urban HAPs and diesel particulate matter, 1996. ....	83
Figure 5-4.	Urban/rural splits by source type for the 1996 national emissions of 188 HAPs. ....	84
Figure 5-5.	Urban/rural splits by source type for the 1996 national emissions of 33 urban HAPs. ....	84
Figure 5-6.	Change in national air toxics emissions – baseline (1990–1993) to 1996. ....	85
Figure 5-7.	Locations for urban and rural air toxics monitors with long-term data. ....	86
Figure 5-8a.	National trend in annual average benzene concentrations in metropolitan areas, 1994–1999. ....	88
Figure 5-8b.	National trend in annual average 1,3-butadiene concentrations in metropolitan areas, 1994–1999. ....	88
Figure 5-8c.	National trend in annual average total suspended lead concentrations in metropolitan areas, 1994–1999. ....	89
Figure 5-8d.	National trend in annual average perchloroethylene concentrations in metropolitan areas, 1994–1999. ....	89
Figure 5-8e.	National trend in annual average styrene concentrations in metropolitan areas, 1994–1999. ....	90
Figure 5-8f.	National trend in annual average toluene concentrations in metropolitan areas, 1994–1999. ....	90
Figure 5-9a.	Trend in annual average benzene concentrations for metropolitan sites in California, 1990–1999. ....	92

Figure 5-9b.	Trend in annual average 1,3-butadiene concentrations for metropolitan sites in California, 1990–1999. ....	92
Figure 5-9c.	Trend in annual average total suspended lead concentrations for metropolitan sites in California, 1990–1999. ....	93
Figure 5-9d.	Trend in annual average perchloroethylene concentrations for metropolitan sites in California, 1990–1999. ....	93
Figure 5-9e.	Trend in annual average styrene concentrations for metropolitan sites in California, 1990–1999. ....	94
Figure 5-9f.	Trend in annual average toluene concentrations for metropolitan sites in California, 1990–1999. ....	94
Figure 6-1.	IMPROVE sites meeting data completeness requirements for sites operating in 1999. ....	100
Figure 6-2.	Comparison of the three visibility metrics (extinction, deciview and visual image). ....	101
Figure 6-2a.	Images of Shenandoah National Park and Yosemite National Park. ....	101
Figure 6-3.	Shenandoah National Park on clear and hazy days and the effect of adding 10 µg/m <sup>3</sup> of fine particles to each. ....	102
Figure 6-4.	Long-term trends for 75th percentile light extinction coefficient from airport visual data (July–September). ....	103
Figure 6-5a.	Visibility trends for 10 eastern U.S. Class I areas for clearest, middle, and haziest 20 percent days in the distribution, 1992–1999. ....	104
Figure 6-5b.	Visibility trends for 26 western U.S. Class I areas for clearest, middle, and haziest 20 percent days in the distribution, 1992–1999. ....	104
Figure 6-6a.	Aerosol light extinction in 10 eastern Class I areas for the clearest 20 percent of the days in the distribution, 1992–1999. ....	105
Figure 6-6b.	Aerosol light extinction in 10 eastern Class I areas for the middle 20 percent of the days in the distribution, 1992–1999. ....	105
Figure 6-6c.	Aerosol light extinction in 10 eastern Class I areas for the haziest 20 percent of the days in the distribution, 1992–1999. ....	105
Figure 6-6d.	Aerosol light extinction in 26 western Class I areas for the clearest 20 percent of the days in the distribution, 1990–1999. ....	106
Figure 6-6e.	Aerosol light extinction in 26 western Class I areas for the middle 20 percent of the days in the distribution, 1990–1999. ....	106
Figure 6-6f.	Aerosol light extinction in 26 western Class I areas for the haziest 20 percent of the days in the distribution, 1990–1999. ....	106
Figure 6-7a.	Class I area significant trends in deciviews for the clearest 20 percent, middle 20 percent, and haziest 20 percent days as summarized in Table 6-1. ....	108
Figure 6-7b.	Class I area significant trends light extinction due to sulfate for the clearest 20 percent, middle 20 percent, and haziest 20 percent days as summarized in Table 6-1. ....	108
Figure 6-7c.	Class I area significant trends for light extinction due to organic carbon for the clearest 20 percent, middle 20 percent, and haziest 20 percent days as summarized in Table 6-1. ....	109
Figure 6-8a.	Aerosol light extinction in (Mm-1) for the clearest 20 percent days and contribution by individual particulate matter constituents, based on 1997–1999 IMPROVE data. ....	110
Figure 6-8b.	Aerosol light extinction in (Mm-1) for the middle 20 percent days and contribution by individual particulate matter constituents, based on 1997–1999 IMPROVE data. ....	110
Figure 6-8c.	Aerosol light extinction in (Mm-1) for the haziest 20 percent days and contribution by individual particulate matter constituents, based on 1997–1999 IMPROVE data. ....	111
Figure 6-9a.	Current visibility impairment expressed in deciviews for the clearest 20 percent days based on 1997–1999 IMPROVE data. ....	112
Figure 6-9b.	Current visibility impairment expressed in deciviews for the middle 20 percent days based on 1997–1999 IMPROVE data. ....	112
Figure 6-9c.	Current visibility impairment expressed in deciviews for the haziest 20 percent days based on 1997–1999 IMPROVE data. ....	113
Figure 7-1.	The National Atmospheric Deposition Program/National Trends Network. ....	116
Figure 7-2.	Annual mean sulfate deposition from precipitation, 1990–1992 vs. 1997–1999. ....	117
Figure 7-3.	Annual mean ammonium deposition from precipitation, 1990–1992 vs. 1997–1999. ....	118
Figure 7-4.	Annual mean nitrate deposition from precipitation, 1990–1992 vs. 1997–1999. ....	119
Figure 7-5.	Rural annual mean SO <sub>2</sub> concentrations from CASTNet, 1990–1992 vs. 1997–1999. ....	120
Figure 7-6.	Rural annual average sulfate concentrations from CASTNet, 1990–1992 vs. 1997–1999. ....	121
Figure 7-7.	CASTNet and subset of 34 long-term monitoring sites used for 1990–1999 trends analysis. ....	121
Figure 7-8.	Trend in ambient sulfates in the rural eastern United States, based on CASTNet monitoring data, 1990–1999. ....	122
Figure 7-9.	Trend in ambient sulfur dioxide in the rural eastern United States, based on CASTNet monitoring data, 1990–1999. ....	122
Figure 7-10.	Rural annual mean ammonium concentrations from CASTNet, 1990–1992 vs. 1997–1999. ....	123
Figure 7-11.	Rural annual mean total nitrate concentrations from CASTNet, 1990–1992 vs. 1997–1999. ....	123
Figure 7-12.	Trend in annual mean ambient sulfur dioxide and sulfate concentrations, based on CASTNet monitoring data, and regional SO <sub>2</sub> emissions from electric utilities in rural eastern United States, 1990–1999. ....	124

Figures 7-13a. and Figure 7-13b. Trend in annual mean ambient sulfur dioxide and sulfate concentrations, based on CASTNet monitoring data, and regional SO <sub>2</sub> emissions from electric utilities in rural eastern United States by calendar quarter, 1990–1999. ....	124
Figures 7-13c. and 7-13d. Trend in annual mean ambient sulfur dioxide and sulfate concentrations, based on CASTNet monitoring data, and regional SO <sub>2</sub> emissions from electric utilities in rural eastern United States by calendar quarter, 1990–1999. ....	125
Figure 7-14. Wet and dry components of sulfur deposition, 1999. ....	126
Figure 7-15. Wet and dry components of nitrogen deposition, 1999. ....	126
Figure A-1. (Multiple NA areas within a larger NA area) Two SO <sub>2</sub> areas inside the Pittsburgh-Beaver Valley ozone NA. ....	226
Figure A-2. (Overlapping NA areas) Searles Valley PM <sub>10</sub> NA partially overlaps the San Joaquin Valley ozone NA. ....	226
Figure B-1. Carbon monoxide monitoring network, 1999. ....	230
Figure B-2. Lead monitoring network, 1999. ....	230
Figure B-3. Nitrogen dioxide monitoring network, 1999. ....	231
Figure B-4. Ozone monitoring network, 1999. ....	231
Figure B-5. PM <sub>10</sub> monitoring network, 1999. ....	232
Figure B-6. Sulfur dioxide monitoring network, 1999. ....	232
Figure B-7. Class I Areas in the Improve Network meeting data completeness criteria. ....	234

# Tables

Table 2-1.	NAAQS in effect as of December 1999 .....	9
Table 2-2.	Milestones in Motor Vehicle Emissions Control .....	14
Table 2-3.	Summary of Changes in Summer 6-9 a.m. Mean Concentrations of NO <sub>x</sub> and TNMOC at PAMS Sites .....	34
Table 2-4.	Biogenic Sources of VOC Emissions By Region .....	37
Table 2-5.	Percent Contribution to PM <sub>2.5</sub> by Component, 1999 .....	52
Table 2-6.	Total SO <sub>2</sub> Emissions from Phase I and Non-Phase I Acid Rain Sources: 1990–1999 (million tons). ....	64
Table 3-1.	Summary of MSA Trend Analyses by Pollutant, 1990–1999 .....	70
Table 3-2.	AQI Categories, Colors, and Ranges .....	72
Table 4-1.	Areas Redesignated Between September 1999 and September 2000 .....	77
Table 5-1.	List of 33 Urban Air Toxics Strategy HAPs .....	81
Table 5-2.	National Summary of Ambient HAP Concentration Trends in Metropolitan Areas, 1994–1999 .....	87
Table 5-3.	National Summary of Ambient HAP Concentration Trends in Rural Areas, 1994–1999 .....	95
Table 6-1.	Summary of Class I Area Trend1 Analysis .....	107
Table A-1a.	National Air Quality Trends Statistics for Criteria Pollutants, 1980–1989 .....	128
Table A-1b.	National Air Quality Trends Statistics for Criteria Pollutants, 1990–1999 .....	130
Table A-2.	National Carbon Monoxide Emissions Estimates, 1970, 1975, 1980, 1985, 1989–1999 (thousand short tons) .....	132
Table A-3.	National Lead Emissions Estimates, 1970, 1975, 1980, 1985, 1989–1999 (short tons) .....	136
Table A-4.	National Nitrogen Oxides Emissions Estimates, 1970, 1975, 1980, 1985, 1989–1999 (thousand short tons) .....	138
Table A-5.	National Volatile Organic Compounds Emissions Estimates, 1970, 1975, 1980, 1985, 1989–1999 (thousand short tons) .....	142
Table A-6.	National PM <sub>10</sub> Emissions Estimates, 1970, 1975, 1980, 1985, 1989–1999 (thousand short tons) .....	148
Table A-7.	Miscellaneous and Natural PM <sub>10</sub> Emissions Estimates, 1970, 1975, 1980, 1985, 1989–1999 (thousand short tons) .....	152
Table A-8.	National Sulfur Dioxide Emissions Estimates, 1970, 1975, 1980, 1985, 1989–1999 (thousand short tons) .....	153
Table A-9.	National PM <sub>2.5</sub> Emissions Estimates, 1990–1999 (thousand short tons) .....	156
Table A-10.	National Ammonia Emissions Estimates, 1990–1999 (thousand short tons) .....	160
Table A-11.	National Long-Term Air Quality Trends, 1980–1999 .....	162
Table A-12a.	National Air Quality Trends by Monitoring Location, 1980–1989 .....	163
Table A-12b.	National Air Quality Trends by Monitoring Location, 1990–1999 .....	164
Table A-13a.	National Air Quality Trends Statistics by EPA Region, 1980–1989 .....	165
Table A-13b.	National Air Quality Trends Statistics by EPA Region, 1990–1999 .....	167
Table A-14.	Maximum Air Quality Concentrations by County, 1999 .....	169
Table A-15.	Maximum Air Quality Concentrations by Metropolitan Statistical Area, 1999 .....	188
Table A-16.	Metropolitan Statistical Area Air Quality Trends, 1990–1999 .....	194
Table A-17.	Number of Days with AQI Values Greater Than 100 at Trend Sites, 1990–1999, and All Sites in 1999 .....	219
Table A-18.	(Ozone only) Number of Days with AQI Values Greater Than 100 at Trend Sites, 1990–1999, and All Sites in 1999 .....	221
Table A-19.	Condensed Nonattainment Areas List(a) .....	223
Table A-20.	Trend in 8-hr ozone concentrations (ppm) exceedances at National Park and National Monument sites, 1990–1999 .....	227
Table A-21.	Onroad and Nonroad Emissions of 21 Mobile Source Air Toxics, 1996 .....	228
Table B-1.	Number of Ambient Monitors Reporting Data to AIRS .....	229

# Acronyms

AIRS	Aerometric Information Retrieval System	NARSTO	North American Research Strategy for Tropospheric Ozone
AQRV	Air-Quality Related Values	NESCAUM	Northeast States for Coordinated Air Use Management
AIRMoN	Atmospheric Integrated Assessment Monitoring Network	NLEV	National Low Emission Vehicle
CAA	Clean Air Act	NMOC	Non-Methane Organic Compound
CAAA	Clean Air Act Amendments	NO <sub>2</sub>	Nitrogen Dioxide
CARB	California Air Resources Board	NO <sub>x</sub>	Nitrogen Oxides
CASAC	Clean Air Scientific Advisory Committee	NPS	National Park Service
CASTNet	Clean Air Status and Trends Network	NTI	National Toxics Inventory
CEMs	Continuous Emissions Monitors	O <sub>3</sub>	Ozone
CFR	Code of Federal Regulations	OTAG	The Ozone Transport Assessment Group
CO	Carbon Monoxide	PAHs	Polyaromatic Hydrocarbons
CMSA	Consolidated Metropolitan Statistical Area	PAMS	Photochemical Assessment Monitoring Stations
DST	Daylight Savings Time	PAN	Peroxyacetyl Nitrate
EPA	Environmental Protection Agency	Pb	Lead
FRM	Federal Reference Method	PBTs	Persistent and Bioaccumulative Toxics
GDP	Gross Domestic Product	PCBs	Polychlorinated Biphenyls
GLM	General Linear Model	PM <sub>10</sub>	Particulate Matter of 10 micrometers in diameter or less
HAPs	Hazardous Air Pollutants	PM <sub>2.5</sub>	Particulate Matter of 2.5 micrometers in diameter or less
IADN	Integrated Atmospheric Deposition Network	POM	Polycyclic Organic Matter
I/M	Inspection and Maintenance Programs	ppm	Parts Per Million
IMPROVE	Interagency Monitoring of PROtected Environments	PSI	Pollutant Standards Index
MACT	Maximum Achievable Control Technology	RFG	Reformulated Gasoline
MARAMA	Mid-Atlantic Regional Air Management Association	RVP	Reid Vapor Pressure
MDN	Mercury Deposition Network	SLAMS	State and Local Air Monitoring Stations
MSA	Metropolitan Statistical Area	SNMOC	Speciated Non-Methane Organic Compound
MDL	Minimum Detectable Level	SO <sub>2</sub>	Sulfur Dioxide
NAAQS	National Ambient Air Quality Standards	SO <sub>x</sub>	Sulfur Oxides
NADP/NTN	National Atmospheric Deposition Program/National Trends Network	TNMOC	Total Non-Methane Organic Compound
NAMS	National Air Monitoring Stations	TRI	Toxic Release Inventory
NAPAP	National Acid Precipitation Assessment Program	TSP	Total Suspended Particulate
		UATMP	Urban Air Toxics Monitoring Program
		VMT	Vehicle Miles Traveled
		VOCs	Volatile Organic Compounds
		µg/m <sup>3</sup>	Micrograms Per Cubic Meter